

CRITICAL BEHAVIOR OF THE ANTIFERROMAGNETIC HEISENBERG MODEL ON A STACKED TRIANGULAR LATTICE

T. Bhattacharya*, A. Billoire, R. Lacaze[†] and Th. Jolicœur[‡]
Service de Physique Théorique de Saclay[‡]

September 24, 1993

We estimate, using a large-scale Monte Carlo simulation, the critical exponents of the antiferromagnetic Heisenberg model on a stacked triangular lattice. We obtain the following estimates: $\gamma/\nu = 2.011 \pm .014$, $\nu = .585 \pm .009$. These results contradict a perturbative $2 + \epsilon$ Renormalization Group calculation that points to Wilson-Fisher $O(4)$ behaviour. While these results may be coherent with $4 - \epsilon$ results from Landau-Ginzburg analysis, they show the existence of an unexpectedly rich structure of the Renormalization Group flow as a function of the dimensionality and the number of components of the order parameter.

SPhT-93/015

cond-mat/9302019

Submitted to: Europhysics Letters

*Present address: MS B285, Group T-8, Los Alamos National Laboratory, NM 87544

[†]Chercheur CNRS

[‡]Laboratoire de la Direction des Sciences de la Matière du CEA.

1 INTRODUCTION

There is at present a satisfactory understanding of the critical behaviour of physical systems where the rotation symmetry group $O(N)$ is broken down to $O(N-1)$ at low temperatures. Several theoretical tools are available to estimate the critical exponents and there is good agreement between these estimates. The Wilson-Fisher fixed point which describes the critical physics can be smoothly followed between two and four dimensions: for $N \geq 3$ the $2+\epsilon$ and $4-\epsilon$ renormalization group expansion merge in a continuous manner.

The situation is much more complicated when the rotation symmetry is *fully* broken in the low-temperature phase. A prominent example is found in the so-called helimagnetic systems where Heisenberg spins are in a spiral arrangement below the critical temperature. It is an interesting question, both theoretically and experimentally, to know the corresponding universality class. A widely studied prototypical model is the antiferromagnetic Heisenberg model on a stacked triangular lattice, which is simple and display commensurate helimagnetic order below a transition point T_c . There is little consensus in the literature regarding critical phenomena associated with this model [1].

This topic has been investigated by use of a $D = 4 - \epsilon$ renormalization group calculation[2]. The corresponding Ginzburg-Landau theory for a N -component vector model involves two N -component bosonic fields. It is found that for N large enough the transition is second order and not governed by the Heisenberg $O(2N)$ Wilson-Fisher fixed point but by a different fixed point which is also non-trivial for $D < 4$. For smaller N , this new fixed point disappears and there is no stable fixed point which is an indication for a fluctuation-induced first-order transition. The dividing universal line between second-order and first-order behaviour is found to be $N_c(D) = 21.8 - 23.4\epsilon + O(\epsilon^2)$. The rapid variation of N_c leaves us rather uncertain about the fate of the case $D = 3$. Clearly more information is needed about the $N_c(D)$ line in the (number of components-dimension)-plane.

A $D = 2 + \epsilon$ renormalization group study has been performed for a system of Heisenberg spins[3] by use of a non-linear sigma model defined on a homogeneous non-symmetric coset space $O(3) \times O(2)/O(2)$. It was found that near two dimensions the system undergoes a second order transition which is governed by the $O(4)$ usual Wilson-Fisher fixed point. In fact the symmetry $O(3) \times O(2)/O(2)$ is dynamically enlarged at the critical point to $O(3) \times$

$O(3)/O(3)$ and $O(3) \times O(3)$ is $O(4)$. This mismatch between the expansions near four dimensions and near two dimensions is quite unusual and does not happen in the well-studied $O(N) \rightarrow O(N-1)$ critical phenomena. As a consequence, the $D = 3$ case remains elusive and a direct study in three dimensions is called for.

Some preliminary Monte Carlo (MC) simulations have shown evidence [4, 5, 6, 7] for a continuous transition in the case of the Heisenberg anti-ferromagnet on a stacked triangular lattice. The exponents found are not compatible with those of the $O(4)$ vector model in three dimensions. We did a large-scale simulation with much better statistics than previous attempts. As a consequence we are able to pin down the transition temperature in a very precise manner and to obtain reliable estimates of critical exponents.

We thus focus on the classical spin model defined by the classical Heisenberg Hamiltonian:

$$H = \sum_{\langle ij \rangle} J_{ij} \mathbf{S}_i \cdot \mathbf{S}_j. \quad (1)$$

The exchange interaction J_{ij} is nonzero ($J_{ij} = 1$ in what follows) between nearest-neighbors of a stacked triangular lattice and the spins are three-component unit vectors. The classical ground state is found by minimizing the Fourier transform $J(\mathbf{Q})$ of the couplings J_{ij} . The spins adopt a planar arrangement on a three-sublattice structure with relative angles 120 degrees.

2 The simulation

We made two successive sets of Monte Carlo simulations of this model, using slightly asymmetric lattices (with periodic boundary conditions) of shape $L^2 L_z$, where L is the linear size inside the planes, and $L_z = 2/3L$ the stacking size.

The first set of simulations was run on a CM-2 8K massively parallel computer. We made runs on lattices of size $12^2 \times 8$ (with a total of 600 000 Monte Carlo sweeps of the lattice), $24^2 \times 16$ (with a total of 1 500 000 sweeps) and $48^2 \times 32$ (with a total of 1 570 000 sweeps). In the first case we used the Heat-Bath algorithm, in the later two cases we used an Hybrid Over-relaxation algorithm[9] where each Heat-Bath sweep is followed by an energy conserving sweep. The next set of simulations was run on CRAY vector

computers on lattices of size $18^2 12$, $24^2 16$, $30^2 20$, $36^2 24$ and $48^2 32$ with 4 to $8 \cdot 10^6$ Hybrid Overrelaxation sweeps. Both simulations concentrate in the immediate vicinity of the transition.

Let us note by $\mathbf{S}_a$ the total spin per site for sublattice a ($a \in [1, 2, 3]$), we measured the magnetization:

$$M_L(\beta) = \frac{1}{3} \sum_a \langle |\mathbf{S}_a| \rangle, \quad (2)$$

the susceptibility:

$$\chi_L(\beta) = \frac{L^2 L_z}{3} \left(\sum_a \langle \mathbf{S}_a^2 \rangle - \langle |\mathbf{S}_a| \rangle^2 \right), \quad (3)$$

and the fourth-order cumulant:

$$B_L(\beta) = 1 - \frac{1}{3} \sum_a \frac{\langle \mathbf{S}_a^4 \rangle}{\langle \mathbf{S}_a^2 \rangle^2}. \quad (4)$$

Our strategy to extract the critical exponents was first to estimate the value of the critical temperature β_c , as the point where $B_L(\beta)$ is L independent and then to estimate the exponents from the following set of equations:

$$\chi_L(\beta_c) \sim L^{\gamma/\nu}, \quad (5)$$

$$M_L(\beta_c) \sim L^{-\beta/\nu}, \quad (6)$$

$$\left. \frac{\partial B_L(\beta)}{\partial \beta} \right|_{\beta=\beta_c} \sim L^{1/\nu} \quad (7)$$

$$\left. \frac{\partial}{\partial \beta} \ln M_L(\beta) \right|_{\beta=\beta_c} \sim L^{1/\nu}. \quad (8)$$

The extrapolation, from the β value used for the simulation, to β_c is done using the Ferrenberg-Swendsen technique [10]. This technique is invaluable to extrapolate in a neighborhood of size $\sim 1/L^{1/\nu}$ (We understand that, when used blindly outside such a tight range, it may give wrong results). An alternative strategy would be to evaluate exponents from the finite size scaling of the maxima of $\chi_L(\beta)$, $M_L(\beta)$, $\frac{\partial}{\partial \beta} B_L(\beta)$ and $\frac{\partial}{\partial \beta} \ln M_L(\beta)$. It would however require to perform simulations at four β values, close to the four (L dependent) points where this quantities reach their maxima, otherwise

accuracy would be lost. The statistical analysis is done with respect to 20 bins, using first-order bias-corrected jackknife (see e.g. Ref.[11] and references therein). The first 20 % of each run is discarded for thermalization.

From simulations of lattices of increasing sizes $L_1 < L_2 < \dots$ converging estimators of β_c are obtained by solving the equations:

$$B_{L_{i-1}}(\tilde{\beta}) = B_{L_i}(\tilde{\beta}). \quad (9)$$

	β_c	γ/ν	$D - 2\beta/\nu$	ν	ν
24-12	1.04401 (38)	2.009 (14) (04)	2.030 (07) (07)	.651 (22) (03)	.604 (10) (01)
48-24	1.04462 (21)	2.033 (18) (14)	1.998 (09) (25)	.590 (20) (04)	.576 (10) (04)
24-18	1.04452 (40)	2.010 (26) (06)	2.011 (08) (10)	.605 (30) (03)	.608 (17) (02)
30-24	1.04388 (36)	1.994 (36) (11)	2.021 (13) (16)	.578 (35) (03)	.583 (19) (03)
36-30	1.04408 (37)	1.988 (55) (16)	2.020 (18) (21)	.605 (55) (12)	.607 (30) (04)
42-36	1.04485 (25)	2.083 (68) (14)	1.984 (20) (31)	.528 (48) (06)	.548 (28) (06)
48-42	1.04427 (30)	1.986 (61) (18)	2.004 (22) (39)	.639 (69) (05)	.577 (27) (03)
48-18	1.04432 (06)	2.011 (07) (12)	2.010 (03) (21)	.590 (09) (06)	.588 (04) (03)
all	N/A	2.011 (07) (12)	2.011 (02) (19)	.585 (07) (06)	.585 (04) (04)

Table 1: Critical temperature and critical exponents estimates, using equations 9, 5, 6, 7 and 8 respectively. First two lines are CM-2 data, next lines the CRAY data. The first column gives the linear sizes of the lattices used. The two numbers inside parenthesis are the estimated statistical errors, direct and induced, on the last digits of the number on their left.

The results can be found in the second column of Tab.1. Within our statistical accuracy all estimates are compatible, there is neither any clear lattice size dependence, nor any discrepancy between the two sets of runs. Our final estimate is (the quoted error is deliberately on the safe side)

$$\beta_c = 1.0443 \pm .0002. \quad (10)$$

We use this value to compute $\chi_L(\beta_c)$, $M_L(\beta_c)$, $\frac{\partial}{\partial\beta}B_L(\beta)$ and $\frac{\partial}{\partial\beta}\ln M_L(\beta)$ in order to estimate γ/ν , β/ν and ν . Results can be found in Tab.1. We quote separately the estimated “direct” statistical errors computed from the dispersion of the results from the 20 bins, and the errors induced by the

uncertainty on the determination of β_c , computed as the (absolute value of the) difference between the values obtained using our best estimate of β_c and the values using the one standard deviation estimate from Eq.10. The last line in Tab.1 gives the results of linear fits of the CRAY data for $\ln(\chi_L(\beta_c))$, $\ln(M_L(\beta_c))$, $\ln(\frac{\partial}{\partial\beta}B_L(\beta)\Big|_{\beta=\beta_c})$ and $\ln(\frac{\partial}{\partial\beta}\ln M_L(\beta)\Big|_{\beta=\beta_c})$ respectively. The results of the fits are stable against omitting the smallest lattice data. The χ^2 is equal to 1.5, 3.7, 1.9 and 4.5 respectively (we expect $\chi^2 \approx 4$). Such values mean that linear fits give good representation of the data for $L \geq 18$ (with the current accuracy).

Our final numbers are $\gamma/\nu = 2.011 \pm .014$, $D - 2\beta/\nu = 2.011 \pm .019$, $\nu = .585 \pm .009$. We observe that hyperscaling is verified within errors, and η is very small ($|\eta| \approx .01$). The value for ν is clearly not compatible with the value for the $O(4)$ fixed point[13] ($\nu \sim .75$).

Published Monte Carlo results for ν are $.53 \pm .02$ [4], $.53 \pm .03$ [5], $.55 \pm .03$ [6] and $.59 \pm .02$ [7]. Our results are in good agreement with the most recent results (and with the result $.57 \pm .02$ from a model of commensurate Heisenberg helimagnet[8]). The methods of analysis are quite different. Ref.[7] uses data taken in a wide region around the transition point and adjust the values of the critical exponents using the old fashioned “data collapsing” method. This method has the disadvantage to give much weight to points with large value of $(\beta - \beta_c)L^{1/\nu}$. We extract the exponents directly from data reweighted to β_c . We have also a much higher statistics: we use 8×10^6 heat-bath + energy-conserving sweeps whereas [7] uses 6-20 times 20000 sweeps of a less efficient algorithm. Our estimated statistical error on the determination of β_c is one order of magnitude smaller than the one in ref.[7].

3 Conclusion

Let us first summarize the knowledge obtained from perturbative RG studies as a function of the dimensionality and the number of components of the order parameter.

- i) There is a universal line $N_c(D)$ separating a first-order region from a second-order region near $D = 4 - \epsilon$. Its slope is known from RG studies[2] as well as the critical value N_c for $D = 4$.
- ii) Large-N studies[12] indicate that the stable fixed point found above $N_c(D)$

persists smoothly in the region $N = \infty$ and $2 < D < 4$.

iii) Smoothness along the $D = 2$ vertical axis can be shown by studying the nonlinear sigma model suited to the N-vector model: it is built on the homogeneous space $O(N) \times O(2)/O(N-2) \times O(2)$ [14]. One finds a single stable fixed point for *all* values of $N \geq 3$. In the large-N limit the exponents from this sigma model are the same as those of the linear model.

If we believe in the perturbative RG results, then necessarily the universal line $N_c(D)$ can only intersect the horizontal axis $N = 3$ between $D=2$ and $D=4$. The simplest hypothesis is then that the plane (N, D) is divided in two regions by the line $N_c(D)$. This line may intersect the $N = 3$ axis at a critical dimension D_c . Our Monte-Carlo results thus imply that D_c is between three and four dimensions since we observe a continuous transition at $D=3$. A possibility would be that the critical behaviour we observe is described by the fixed point found in the neighborhood of $D=4$.

However one has to note that the sigma model approach shows that this fixed point is $O(4)$ for $N=3$. This is clearly ruled out by our data. The simplest scenario suggested by i-iii above is wrong. It may be that the whole sigma model approach breaks down since perturbative treatment of a sigma model neglects global aspects. Previous studies indeed have performed, as usual, only perturbation theory for spin-wave excitations[3]. One may invoke the peculiar vortices present in the model[16] since $\Pi_1(SO(3)) = Z_2$ that form loops in $D=3$ as responsible of the breakdown of the sigma model. However vortices are also excluded from the perturbative calculations near $D=4$ and thus it would be difficult in this case to believe in what is found perturbatively. It may be also that the fixed point found in $2 + \epsilon$ expansion is destabilized by some operators containing high powers of gradients [15].

We have thus obtained the critical behaviour of a magnet with a canted ground-state. This calls for a deeper understanding of the Renormalization Group behaviour of canted systems since there is no obvious relationship between the different perturbative approaches, contrary to the case of collinear vector magnets. We note for the future that it would be interesting to obtain the critical behaviour of the XY canted systems, much more relevant to the experimental situation.

Acknowledgements


We would like to thank Guy Decaudain, David Lloyd Owen, Gyan Bhanot and Roch Bourbonnais of TMC for their support. We also thank H. T. Diep, A. P. Young and D. P. Belanger for various informations about this subject.

References

- [1] For a review see: M. L. Plumer and A. Caillé, J. Appl. Phys. **70**, 5961 (1991).
- [2] T. Garel and P. Pfeuty, J. Phys. C**9**, L245 (1976); D. Bailin, A. Love and M. A. Moore, J. Phys. C**10**, 1159 (1977).
- [3] P. Azaria, B. Delamotte and Th. Jolicœur, Phys. Rev. Lett. **64**, 3175 (1990).
- [4] H. Kawamura, J. Phys. Soc. Jpn **54**, 3220 (1985).
- [5] H. Kawamura, J. Phys. Soc. Jpn **56**, 474 (1986).
- [6] H. Kawamura, J. Phys. Soc. Jpn **58**, 584 (1989).
- [7] H. Kawamura, J. Phys. Soc. Jpn **61**, 1299 (1992).
- [8] H. T. Diep, Phys. Rev. **B 39**, 397 (1989).
- [9] A.D. Kennedy, Proceedings of the International Symposium on Lattice Field Theory LATTICE 92, September 15-19 1992 Amsterdam.
- [10] A.M. Ferrenberg and R.H. Swendsen, Phys. Rev. Lett. **61** 2635 (1988);
See also A.M. Ferrenberg and R.H. Swendsen, Phys. Rev. Lett. **63** 1195 (1989).
- [11] A. Billoire, R. Lacaze, and A. Morel, Nucl. Phys. B **370** 773 (1992).
- [12] H. Kawamura, Phys. Rev. **B38**, 4916 (1988).
- [13] H. E. Stanley in “Phase transitions and critical Phenomena”, Vol. III, C. Domb and M. S. Green eds., Academic Press, New York, 1974; J.C. Le Guillou and J. Zinn-Justin have also obtained unpublished estimates of the $O(4)$ exponents by field-theoretic methods that are very close to the series estimates.

- [14] P. Azaria, B. Delamotte, F. Delduc and Th. Jolicœur, Saclay preprint SPhT/93-044, to appear in Nuclear Physics B[FS].
- [15] G. E. Castilla and S. Chakravarty, Phys. Rev. Lett. **71**, 384 (1993).
- [16] H. Kawamura and S. Miyashita, J. Phys. Soc. Jpn **53**, 4138 (1984).

NUMBER OF COMPONENTS



DIMENSION

